

E-cadherin Monoclonal Antibody (ECCD-2)

Product Details

Size	100 µg
Species reactivity	Mouse
Published species	Avian, Dog, Fish, Hamster, Human, Mouse, Pig, Rat, Tag
Host / Isotype	Rat / IgG2a
Class	Monoclonal
Type	Antibody
Clone	ECCD-2
Conjugate	Unconjugated
Immunogen	Mouse liver E-cadherin fragment.
Form	Lyophilized
Concentration	2.0 mg/mL
Purification	Ion-exchange chromatography
Storage buffer	TBS, pH 7.5, with 1% BSA
Contains	no preservative
Storage conditions	4°C
RRID	AB_2533005

Applications	Tested Dilution	Publications
Flow Cytometry (Flow)	-	12 Publications
Immunocytochemistry (ICC/IF)	-	59 Publications
Immunohistochemistry (IHC)	-	95 Publications
Immunohistochemistry (Frozen) (IHC (F))	10 µg/mL	12 Publications
Immunohistochemistry (Paraffin) (IHC (P))	-	11 Publications
Immunohistochemistry (PFA fixed) (IHC (PFA))	-	2 Publications
Immunomicroscopy (IM)	-	1 Publication
Immunoprecipitation (IP)	-	2 Publications
Miscellaneous Pubmed (Misc)	-	34 Publications
Western Blot (WB)	10 µg/mL	34 Publications

Product Specific Information

Source: Monoclonal antibody was obtained by fusing the P3-X63-Ag8 mouse myeloma cell line with spleen cells of Donryu rat after immunization with mouse teratocarcinoma and mouse liver E-cadherin fragment.

Specificity: This antibody specifically reacts with mouse E-cadherin. This antibody does not inhibit E-cadherin-dependent cell-cell contact.

Cross reactivity: This antibody reacts with human and dog E-cadherin. This antibody does not react with bovine or chicken E-cadherin.

Western blot can be performed under reducing or non-reducing conditions.

Reconstitution and dilution: dissolve the lyophilized antibody in 50 µL of distilled water to a final concentration 2.0 mg/mL. This solution can be used as a stock solution. If dilution is necessary for your application, dilute the stock solution in 500 µL or more of the following dilution solution: 20mM TBS (pH 7.5), 10mM CaCl₂, 1.0% BSA, 0.1% Na₃N. Note that TBS should be used for dilution, as PBS causes precipitation.

Storage: When stored at 4°C, 0.1% sodium azide should be added as a preservative. Store the antibody at a minimum concentration of 2.0 mg/mL; a lower antibody concentration may result in decreased stability. The stock solution (2.0mg/mL) should be stored in aliquots at -20°C for 1year, or stored at 4°C for 6 months after adding 0.1% sodium azide. Avoid repeated freeze-thaw cycles. Diluted antibody should not be stored.

Visit thermofisher.com to view references and figures.

Products are under warranty for one year from the date of shipment unless otherwise stated on this document or the product label.

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample.

NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S

EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.