

OxPhos Rodent WB Antibody Cocktail

Product Details

Size	300 µg
Species reactivity	Bovine, Human, Mouse, Non-human primate, Rat
Published species	Human, Mouse, Rat, Sheep
Host / Isotype	Mouse / Ig
Class	Cocktail
Type	Antibody
Clone	Cocktail
Conjugate	Unconjugated
Form	Liquid
Concentration	1.5 mg/mL
Purification	purified
Storage buffer	HEPES buffered saline
Contains	0.02% sodium azide
Storage conditions	4°C
RRID	AB_2533835

Applications

Tested Dilution

Publications

Western Blot (WB)	Assay-dependent	41 Publications
-------------------	-----------------	-----------------

Product Specific Information

This product is a cocktail of multiple OxPhos antibodies.

C-I-20 (ND6) (clone) 20E9DH10C12 - Bovine Complex I

Purification Method: Antigen Affinity Chromatography

C-II-30 (FeS) (clone) 21A11AE7 - Bovine Heart Complex II

Purification Method: IgG Fraction

C-III-Core 2 (clone) 13G12AF12BB11 - Bovine Heart Complex II, III

Purification Method: IgG Fraction

C-IV-I (clone) 1D6E1A8 - Human Complex IV subunit I

Purification Method: Ammonium Sulfate Precipitation

C-V-alpha (clone) 15H4C4 - Bovine Complex V

Purification Method: Antigen Affinity Chromatography

The kit is suitable for Western Blotting analysis of the relative levels of the 5 OxPhos complexes in mitochondrial preparations from mouse, rat, human, or bovine sources. These clones were chosen because they are against a subunit that is labile when its complex is not assembled. Moreover, the combination is readily resolved in SDS-PAGE when the appropriate gel conditions are used. Also, COXI is a highly hydrophobic protein and appears as a broad band at ~35 kDa (not at its true molecular weight at 57 kDa). It is very sensitive to heating. Therefore, the samples, including the positive control, should not be heated over 50°C before loaded on the gel.

The antibody cocktail (1.5 mg/mL) should be diluted 250x to a final working concentration of 6.0 µg/mL for Western blotting. We recommend using PBS with 1% milk as the antibody diluent and using a high pH CAPS/PVDF transfer protocol.

Visit thermofisher.com to view references and figures.

Products are under warranty for one year from the date of shipment unless otherwise stated on this document or the product label.

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample.

NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF: (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.