

Rabbit anti-Goat IgG (H+L) Secondary Antibody, TRITC

Product Details		
Size	1.5 mg	
Species reactivity	Goat	
Host / Isotype	Rabbit / IgG	
Class	Polyclonal	
Туре	Secondary Antibody	
Conjugate	TRITC	
Target class	lgG	
Form	Lyophilized	
Concentration	1.5 mg/mL	
Purification	Affinity chromatography	
Storage buffer	PBS, pH 7.6, with 15mg/mL BSA	
Contains	0.05% sodium azide	
Storage conditions	4° C	
RRID	AB_228396	

Applications	Tested Dilution	Publications
Flow Cytometry (Flow)	Assay-dependent	-
Immunocytochemistry (ICC/IF)	Assay-dependent	-
Immunohistochemistry (IHC)	Assay-dependent	-
Immunoprecipitation (IP)	Assay-dependent	-
Western Blot (WB)	Assay-dependent	-

Product Specific Information

Concentration may vary slightly from lot-to-lot, see lot-specific datasheet for exact concentration.

Antibody Specificity: This antibody reacts with heavy chains on goat IgG as well as with the light chains common to most goat immunoglobulins, based on electrophoresis. No antibody was detected against non-immunoglobulin serum proteins. However, this antibody may cross-react with immunoglobulins from other species. Product # 31650 is recommended for use in Western blot, IF, ICC, IHC, IP and FACS applications. The suggested dilution range for most applications is 1:50-1:200.

Fluorophore: Tetramethyl rhodamine 5 (and 6)-isothiocyanate (TRITC)

Amax = 550 nm; Emax = 570 nm

Fluorophore/Protein Ratio: A550/A280 = 0.50

Restoration and Storage: Store product at 4°C until opened. Restore with 1.1 mL distilled water. Centrifuge product if it is not completely clear after standing for 1-2 hours at room temperature. Product may be stored for up to several weeks at 4°C as an undiluted liquid. After dilution, do not use for more than one day. To extend the shelf-life of this product, add an equal volume of glycerol to make a final concentration of 50% glycerol and store at -20°C; or as an alternative, aliquot without glycerol and store at -80° or below, avoiding repeated freezing and thawing. Country of Origin: USA

Visit thermofisher.com to view references and figures.

Products are under warranty for one year from the date of shipment unless otherwise stated on this document or the product label.

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample.

anu quanty or goous and does not represent that any Product will contom to such model or sample. NO OTHER WARRANTES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT, BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) MIPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless of the Product in the Product on the Product uses, or any type of consumption by or application to human or animals.

1